



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Inbreeding, immune defence and ectoparasite load in different mockingbird populations and species in the Galapagos Islands

Hoeck, Paquita E A ; Keller, Lukas F

Abstract: Inbreeding may impair an individual's immune system, render it more susceptible to disease and hence contribute to the extinction risk of small and isolated populations, as often found on islands. So far, surprisingly few studies have assessed the effects of inbreeding on immunocompetence in wild populations. Using 26 microsatellite loci and genetic data from museum specimens and contemporary samples, we calculated short-term and long-term inbreeding in 13 different mockingbird populations covering the range of all 4 species in the Galapagos Islands and compared them with three different measures of innate immunity and ectoparasite load. We found no significant effect of either measure of inbreeding on natural antibody or complement enzyme titres, heterophil-lymphocyte ratio or feather louse abundance. Hence, our results do not support a link between inbreeding and immunocompetence. However, overall statistical power and repeatabilities of antibody and complement enzyme titres were low. Nevertheless, generally, natural antibody titres were high suggesting that the mockingbirds may be equipped with a strong first line of defence, as found in other island species.

DOI: <https://doi.org/10.1111/j.1600-048X.2012.05725.x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-68811>

Journal Article

Accepted Version

Originally published at:

Hoeck, Paquita E A; Keller, Lukas F (2012). Inbreeding, immune defence and ectoparasite load in different mockingbird populations and species in the Galapagos Islands. *Journal of Avian Biology*, 43(5):423-434.

DOI: <https://doi.org/10.1111/j.1600-048X.2012.05725.x>

TITLE

**Inbreeding, immune defence and ectoparasite load in different mockingbird
populations and species in the Galápagos Islands**

Paquita E. A. Hoeck & Lukas F. Keller

Institute of Evolutionary Biology and Environmental Studies, University of Zurich,
Winterthurerstrasse 190, 8057 Zurich, Switzerland.

Corresponding author:

Paquita E. A. Hoeck

Institute for Conservation Research, San Diego Zoo Global, 15600 San Pasqual Valley Road,
Escondido, CA 92027-7000, USA

Phone: +1-760-291-5439

RUNNING TITLE

Inbreeding and immunity of mockingbirds

ABSTRACT

Inbreeding may impair an individual's immune system, render it more susceptible to disease and hence contribute to the extinction risk of small and isolated populations, as often found on islands. So far, surprisingly few studies have assessed the effects of inbreeding on immunocompetence in wild populations. Using 26 microsatellite loci and genetic data from museum specimens and contemporary samples, we calculated short-term and long-term inbreeding in 13 different mockingbird populations covering the range of all 4 species in the Galápagos Islands and compared them with three different measures of innate immunity and ectoparasite load. We found no significant effect of either measure of inbreeding on natural antibody or complement enzyme titres, heterophil-lymphocyte ratio or feather louse abundance. Hence, our results do not support a link between inbreeding and immunocompetence. However, overall statistical power and repeatabilities of antibody and complement enzyme titres were low. Nevertheless, generally, natural antibody titres were high suggesting that the mockingbirds may be equipped with a strong first line of defence, as found in other island species.

KEYWORDS

Genetic diversity, immunocompetence, birds, lice, natural antibodies

INTRODUCTION

Small and isolated populations are not only more susceptible to demographic and environmental stochastic events (Reed 2004), but also suffer from inbreeding and a faster loss of genetic diversity through genetic drift (Allendorf and Luikart 2007). One way in which inbreeding and loss of genetic diversity may increase extinction risks of small populations is by making them more vulnerable to infectious diseases (e.g. McCallum and Dobson 1995, Smith et al. 2009). Inbred individuals and populations have been shown to exhibit a decrease in parasite or pathogen resistance (e.g. Cassinello et al. 2001, Coltman et al. 1999, Puurtinen et al. 2004, Pearman and Garner 2005, Acevedo-Whitehouse et al. 2003, Hedrick et al. 2001, Ross-Gillespie et al. 2007) or lowered immune response (e.g. Reid et al. 2007, Reid et al. 2003), albeit not universally (e.g. Giese and Hedrick 2003, Gerloff et al. 2003, Sandland et al. 2009). Results are therefore still contradictory and more empirical data are needed to evaluate the overall impacts of inbreeding on the ability of populations to cope with disease under natural environmental conditions (Smith et al. 2009, Keller and Waller 2002, Spielman et al. 2004, Hawley et al. 2010).

Small host populations on isolated islands are particularly susceptible to foreign pathogens (McCallum and Dobson 1995), and diseases have caused dramatic population declines and extinctions of avian species on islands (Van Riper et al. 2002, Van Riper et al. 1986). In the Galápagos Islands, the introduction of alien avian diseases has become a major concern leading to the initiation of a number of avian health survey projects (Gottdenker et al. 2005, Soos et al. 2008, Parker et al. 2006, Wikelski et al. 2004), including a disease survey of the endangered Floreana mockingbird (*Mimus trifasciatus*; Deem et al. 2011). *Mimus trifasciatus* is one of four endemic, allopatrically living mockingbird species found in the Galápagos (Harris 1974) and, since its extinction from the main island of Floreana in the late 1800s, its distribution has been restricted to two small satellite islands

(Fig. 1). In a recent study that covered the range of all four Galápagos mockingbird species (Fig. 1) and many different populations, we detected pronounced differences in average inbreeding levels among species and populations (Hoeck et al. 2010b) rendering these isolated populations of different size an ideal natural system to study the effects of genetic diversity on different immune components. Additionally, the availability of genetic data from historical samples over 100 years old allows differentiating between recent inbreeding vs. historically low levels of genetic diversity (Hoeck et al. 2010b). This is of interest because populations that have undergone historical processes of inbreeding may have successfully purged some of their immunity-related genetic load (Crnokrak and Barrett 2002, Ross-Gillespie et al. 2007), resulting in a weaker association between inbreeding and immunity.

In this study, using genetic data from historic and contemporary samples, we determined levels of genetic diversity and inbreeding at the population level and their effects on immunocompetence, i.e. the ability to prevent or control infections by pathogens and parasites (Norris and Evans 2000). We studied 13 populations covering the range of the four mockingbird species in the Galápagos (Fig. 1) and, using two different measures of F_{st} as estimates of population-level inbreeding, we investigated the relationship between inbreeding and multiple immune parameters. We tested a) the innate immunity by counting different types of white blood cells, b) the innate humoral immune response by assessing natural antibody and complement enzyme activity, and c) determined ectoparasite load to test for a potential relationship between inbreeding and parasite burden.

The avian immune system relies on three major defence mechanisms to resist infection by parasites and disease, namely innate immunity, and humoral and cell-mediated acquired immunity (Cheng and Lamont 1988). All three components have been shown to be under genetic control (Sarker et al. 2000) and could hence be affected by inbreeding. Innate immune defences are constitutive and induced rapidly and therefore most important against first exposures to pathogens and quickly growing infections, resulting in non-specific defence

reactions such as inflammatory reaction or phagocytosis. The typical response to infectious or inflammatory diseases in birds is an increase in the total white blood cell (WBC) count. Increased lymphocyte numbers have been detected in infected individuals (e.g. Ricklefs and Sheldon 2007, Apanius et al. 2000) and high ratios of heterophils to lymphocytes (H:L ratios) in blood samples reliably indicate stress (Davis et al. 2008). Therefore, the heterophil:lymphocyte ratio has been suggested to serve as a measurement of infection status and physiological stress in poultry (e.g. Gross and Siegel 1983, Maxwell 1993) and wild birds (Tompkins et al. 2006). We determined relative abundances of heterophils and lymphocytes to test for a link between inbreeding and H:L ratio.

Although innate immunity mediated by phagocytic and bactericidal WBCs represents a potent first line of defence against pathogens, natural antibodies (NABs) provide a supplementary or alternative constitutive defence (e.g. Ochsenbein and Zinkernagel 2000, Zouali 2001). As they are directly encoded in nuclear DNA (Belperron and Bockenstedt 2001) and levels have been shown to respond to selection on other immune components in chickens (Parmentier et al. 2004), genetic differences may influence their titres. NAB levels may hence covary with genetic diversity, as has been shown for other immunologically important proteins in vertebrates such as MHC (Miller and Lambert 2004). Furthermore, NAB response is hypothesized to predict the strength of the adaptive immune response (Kohler et al. 2003), providing a link between the innate and acquired humoral immunity (Lammers et al. 2004, Ochsenbein and Zinkernagel 2000). Together with the complement enzymes, NABs initiate the complement enzyme cascade which eventually leads to cell lysis (Carroll and Prodeus 1998). In a hemolysis-hemagglutination test, we determined agglutination (which arises from NABs only) and lysis (reflecting an interaction between NABs and complement enzymes) to estimate the strength of the constitutive innate immune system (Matson et al. 2005).

We also tested whether inbreeding correlates with ectoparasite load by determining the abundance of two different louse lineages in the plumage of the mockingbirds (Stefka et al. 2011). Ectoparasites encounter host immune defences when they feed on blood or living tissue (Marshall 1981) and may therefore stimulate a spectrum of different immune responses that potentially impair their development or even kill them (Wikel and Alarcon-Chaidez 2001). A link between genetic diversity and immunity and/or ectoparasite abundance was detected in some cases (Whiteman et al. 2006, Luong et al. 2007), whereas no such correlation was found in others (Taylor and Jamieson 2007).

We made the following predictions: a) If inbreeding negatively affects immunocompetence, e.g. through deleterious mutations or less variable immune components, more inbred populations should show lower levels of average innate and innate-humoral immunity and carry more ectoparasites (i.e. show higher infestation intensity) relative to outbred populations, and b) populations that experienced relatively recent, fast inbreeding are expected to show lower immunocompetence and defence against ectoparasites than populations with a long history of low genetic diversity.

MATERIALS AND METHODS

Sample collection

We visited the Galápagos Islands from December-March in three consecutive years (2006-2008) and collected samples on 11 different islands and at 13 different locations (Fig. 1 & Table 1). On St. Cruz and San Cristóbal, we obtained samples from two different locations, separated by approx. 10 km and 35 km, respectively. We distinguished between the two locations on the same island as previous genetic analyses detected substantial within-island genetic differentiation (Hoeck et al. 2010b). Four islands (Champion, Gardner-

by-Floreana, Española and St. Cruz) were visited in all three years to account for variation between years in the parameters estimated here.

Mockingbirds were captured in potter traps and, in some areas, also in mist nets and bled immediately after capture. Blood samples were obtained by a small puncture of the wing vein collecting between 20-140 μ l of blood (less than 1% of the body weight of a bird) in heparinized microtubes. The blood was transferred immediately into Eppendorf tubes for centrifugation on site. Small drops of blood were used to produce two blood smears and one drop was transferred onto filter paper treated with 0.5 M EDTA for later DNA extraction. Blood smears were air-dried and fixed and stained with Diff-Quick solution (Medion Diagnostics GmbH, Düringen, Switzerland). Whole blood was centrifuged in the field and plasma was transferred into a liquid nitrogen container where it was kept frozen until used for the hemolysis-hemagglutination assay in the lab in Switzerland.

White blood cell counts

Blood smears were examined under 1000 x magnification with oil immersion and the proportion of different types of white blood cells (WBCs), i.e. heterophils, lymphocytes, eosinophils, basophils and monocytes, was assessed by counting a total of 100 WBCs. All counts were performed by the same person (PEAH). To estimate variation within the same smear (i.e. repeatability), we repeated the count of 100 WBCs for 20 different individuals. Additionally, to estimate variation between two smears taken from the same individual at the same time, we also analysed the second smear of 22 individuals. We only used data for lymphocytes and heterophils as the most numerous WBCs in our analyses.

As a higher number of WBCs relative to red blood cells (RBCs) can indicate greater innate immune capacity (Zuk and Johnsen 1998), we also counted the relative abundance of WBCs in a subsample of 10 individuals for each of seven populations sampled in 2007. Relative WBC numbers were determined using a slide scanner (Zeiss Mirax Midi Slide Scanner, Carl Zeiss Inc.) and selecting a field covering approx. 10,000 - 40,000 RBCs. RBCs

were counted automatically with the scanner's software (Pannoramic Viewer 1.14.50), and WBCs in the scanned field were subsequently counted by eye.

Hemolysis-hemagglutination assay

Agglutination and lysis titres were assessed with a hemolysis-hemagglutination assay as described in Matson et al. (2005), with higher scores indicating more effective immune responses. Plasma samples were placed on plates randomly, serially diluted twofold with saline and incubated with rabbit red blood cells (Harlan Laboratories UK Ltd.) for 90 min at 37°C. Because we only obtained small volumes of plasma from most mockingbirds, we omitted the first, undiluted row in the assay, hence working with 25 µl of plasma and a dilution series from 1/2 to 1/4096. A plasma sample from a single chicken individual (Harlan Laboratories UK Ltd.) was used as a control. We determined the dilution step at which either the agglutination or lysis reaction stopped (titre score) and took digital images. Titre scores were confirmed a few days later using the digital images only. All scoring was carried out blindly with respect to the identity of the individual birds and was always performed by the same person (PEAH). Sixteen individuals were tested on two different plates to estimate within-individual repeatability among different assay plates. To account for differences between plates processed at different times, all mockingbird lysis and agglutination scores were corrected for the chicken control sample (that was run at the same time) by subtracting the score of the chicken from the mockingbird scores.

Ectoparasite collection and counting

To quantify ectoparasite load, we dust-ruffled a subset of mockingbirds (Table 1) following the method described in Walther and Clayton (1997) by applying 0.7g of pyrethron powder (non-toxic to birds; 0.3% natural flower-extract pyrethrum and 1% piperonyl butoxid; Vetyl-Chemie GmbH, Germany) to the plumage of the birds (all feather tracts except the head). Dust-ruffling time was kept short to reduce stress for the birds under the hot field

conditions. Dusting was performed for 2.5 min, followed by 1 min of incubation and 2.5 min ruffling over a clean plastic tray to extract ectoparasites which were subsequently stored in 97% ethanol until counted and identified in the lab. Only adult and nymphal lice were considered for further analyses as they were the largest and clearly most abundant ectoparasites detected (77% of all organisms detected in the plumage). Identification of louse genera was done by Vincent Smith, Natural History Museum, UK. Lice belonged to two different genera, *Brueelia galapagensis* (Ischnocera) and *Myrsidea nesomimi* (Amblycera). Because *Brueelia* belongs to the feather-chewing lice, it is unlikely to have a direct effect on the birds' immune system, whereas *Myrsidea* lice may cause an immune reaction through their tissue- and blood-feeding behaviour (Moller and Rozsa 2005). However, because feather-chewing lice are also known to have negative effects on host fitness (Booth et al. 1993, Clayton et al. 1999) and might select on behavioural or physical host defences that may be sensitive to host genetic diversity (Whiteman et al. 2006), we included both *Myrsidea* and *Brueelia* lice in our analyses.

DNA extraction and microsatellite analysis

Extraction of DNA from blood on filter paper and amplification at 26 microsatellite loci (MpAAT18, 25, 26, 45, 83 and 96, and Nes01, 03, 04, 05, 06, 07, 08, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22 and 23) were performed as described previously (Hoeck et al. 2009). Microsatellite loci were amplified in six independent multiplex reactions (Panel A-D, Hoeck et al. 2009; Panel E with MpAAT18, 25, 45 and 83; Panel F with MpAAT26 and 96 under the same PCR conditions as Panel B & C). Fragment analyses were performed on a 3730 DNA Analyser using Gene-Scan-500 LIZ size standard (ABI) and Genemapper v.4 software (ABI) followed by manual proofreading of genotypes.

Because morphology-based sexing is difficult, we performed molecular sexing by amplifying the CHD-W and CHD-Z genes (Griffiths et al. 1998) using redesigned primers (Hoeck et al. 2009).

Genetic diversity and inbreeding estimates

Hardy-Weinberg equilibrium for each locus was tested with allele randomizations within samples (1,000 permutations per test) and overall samples (10,000 permutations) using FSTAT 2.9.3.1 package (Goudet 2001) and adjusting for multiple comparisons using Bonferroni corrections. Genotypic equilibrium between all pairs of loci in each population was tested using G-statistics with Bonferroni corrections and 84,500 permutations in FSTAT. Within population genetic diversity was estimated using standard genetic parameters such as number of alleles (NA), allelic richness (AR, standardized to the smallest sample size) and expected heterozygosity (He) as calculated in FSTAT. As NA and AR correlated strongly with He (NA: $r^2 = 0.89$, $p < 0.001$; AR: $r^2 = 0.94$, $p < 0.001$), we only used He as a measure of population genetic diversity in our analyses.

Inbreeding is always a relative measure and different estimates depend on their population of reference (Keller and Waller 2002). Hence, they also relate to different time periods in the past during which inbreeding could accumulate (Biebach and Keller 2010). In randomly mating populations, overall inbreeding can be expressed as Wright's F_{st} (Keller and Waller 2002). Here, we calculated two different measures of F_{st} to distinguish between two different time scales during which inbreeding accumulated in each population: recent (i.e. short-term) inbreeding and long-term inbreeding. As a measure of recent inbreeding we used the previously published "temporal F_{st} " Table 1 (Hoeck et al. 2010b). This temporal F_{st} quantifies the increase in inbreeding within each island/population over the last century by comparing the genetic diversity of museum specimens collected in 1906 to the one found for the contemporary samples (Hoeck et al. 2010b). The historic data set consisted of 302 specimens from all 11 islands investigated here (sample size: 11-29 specimens/island) that were genotyped at 16 microsatellite loci. These 16 loci were used to quantify changes in microsatellite allele frequencies since 1906, i.e. temporal differentiation (temporal F_{st}) between historic and contemporary samples within each island (Hoeck et al. 2010b). To

quantify long-term inbreeding, we calculated population-specific F_{st} values (Biebach and Keller 2010) using the software 2mod (Ciofi et al. 1999) and selecting the “infer model” mode that we ran with 500,000 iterations. Inbreeding can occur due to both, subdivision in finite populations and non-random mating within subpopulations, and overall inbreeding is given by $(1-F_{it}) = (1 - F_{is})(1 - F_{st})$ (Wright 1969, p. 295). Since we did not detect any inbreeding due to non-random mating, i.e. $F_{is}=0$ (it was not significantly different from zero in any population, see below), overall inbreeding equals Wright’s F_{st} (i.e. $F_{it}=F_{st}$; Keller and Waller 2002). We call this F_{st} “population-specific F_{st} ” because it quantifies total inbreeding that accumulated due to drift (as there is no inbreeding due to non-random mating) relative to the last common ancestral population (Biebach and Keller 2010).

Heterozygosity-heterozygosity correlations

The use of microsatellite markers to predict levels of inbreeding has been questioned (e.g. Balloux et al. 2004, DeWoody and DeWoody 2005, Ljungqvist et al. 2010). We therefore estimated the degree to which heterozygosity is correlated across unlinked markers following the method described in Balloux et al. (2004): assuming that microsatellite and genome-wide heterozygosity are correlated, heterozygosity estimated from one set of microsatellites should be positively correlated with heterozygosity from an independent set of microsatellites from the same individual (‘heterozygosity–heterozygosity correlations’). We randomly split the 26 loci into two sets of 13 independent loci and performed the analysis, a) for each population separately to estimate heterozygosity-heterozygosity correlations on the individual level within a population, and b) in a separate analysis, including all populations and individuals to estimate correlations at the population level, both times using 1,000 iterations to calculate the distribution of the correlation coefficients between the two sets of loci. We detected high correlations when including all individuals and populations in the analysis (mean correlation coefficient: $0.65 \pm SD 0.03$), but low or no correlation when doing analyses for each population separately (correlation coefficients ranging from -0.24 to 0.16 with

distributions including 0). We therefore did not include individual multilocus heterozygosity in any of our statistical models, but restricted our inbreeding – immunocompetence analyses to the population level.

Statistical analyses

Repeatabilities to test for within and among individual variation in heterophil-lymphocyte (H:L) ratios, as well as lysis and agglutination scores were calculated according to Lessells & Boag (1987). To stabilize variances and reach normal distributions, Box-Cox transformations were performed for the response variables H:L ratio ($\lambda = 0$) and number of feather lice ($\lambda = 0.25$; Box and Cox 1964).

Our aim was to assess the relationship between inbreeding and immunocompetence estimates. However, first we carried out an analysis to correct for environmental effects that may also influence our immune and ectoparasite estimates, and could hence lead to differences between islands. To this end, we performed individual-based generalized linear mixed model (GLMM) analyses including all sampled individuals, to account for any sex, year or species effects. We included sex, year and species as fixed effect variables, species nested within island as random effect, and performed separate analyses for each of the four response variables, i.e. agglutination, lysis, H:L ratio and number of feather lice, to obtain the least squares means for these variables. We then used these least squares means to investigate the relationship between levels of inbreeding and immunocompetence. We carried out population-level generalized linear models (GLM) analyses using temporal F_{st} and population-specific F_{st} as fixed effect variables and the least squares means from the individual-based analyses as response variables, again performing a separate analysis for each of the four responses. Qualitatively, using the least squares means instead of the true population means for the four response variables did not provide any differing results (data not shown). Therefore, for ease of interpretation, we used the true population means for agglutination, lysis, H:L ratio and number of feather lice in our final models. We also

performed analyses including mean population H_e as a further explanatory variable, but omitted it in our final analyses because H_e and population-specific F_{st} were highly correlated ($r^2 = 0.96$, $p < 0.0001$) and H_e did not add any additional information to our model.

Not all information was available for all individuals. Sample sizes therefore varied slightly among analyses (Table 1 & 2). All analyses were performed using the software JMP Version 8 and SAS Version 9.1 (SAS Institute Inc., Cary, NC) with significance set to $\alpha = 0.05$.

RESULTS

White blood cell counts

Within-smear repeatability was 82.5% ($F_{19,20} = 12.9$, $p < 0.0001$) and between-smear repeatability 84.7% ($F_{21,22} = 16.4$, $p < 0.0001$), demonstrating that our white blood cell counts were reliable. Mean heterophil:lymphocyte (H:L) ratio varied considerably among islands with the two *M. macdonaldi* populations and the birds from Rábida showing the highest values (Table 1 & Fig. 2a).

Relative numbers of white blood cells (WBCs/RBCs) in blood smears were unrelated to the H:L ratio of individual smears ($r^2 = 0.03$, $p = 0.15$) and also showed no significant correlation with any of the genetic diversity (NA, AR, H_e) or inbreeding (F_{st}) measures (all p -values > 0.46 and r^2 -values < 0.11 , data not shown). Therefore, we did not proceed to determine the WBC/RBC ratio for the remaining samples (i.e. the entire data set).

Hemolysis-hemagglutination assay

Repeat tests of the 16 individuals that were analysed twice (but on different assay plates) showed that repeatability was 8% for the hemagglutination reaction ($F_{15,16} = 1.24$, $p =$

0.33), whereas it was 58% for the hemolysis reaction ($F_{15,15} = 4.57$, $p = 0.003$). Hence, most of the variation in agglutination occurred within individuals.

Agglutination scores of individuals ranged from 3.5 to 11 (the maximum possible) with an overall mean score of 8.8 (\pm SD 1.2) and lysis ranged from 0 to 7 with a mean of 3.4 (\pm SD 1.2). Overall, lysis and agglutination scores varied among populations (Fig. 2b & 2c), with a difference of nearly two scores in mean agglutination titres and 1.5 scores in mean lysis titres (Table 1). Interestingly, no lysis occurred in any of the samples collected on Rábida.

Ectoparasites

We dust-ruffled 196 individuals from all 13 populations (Table 1) and collected a total of 1,482 *Myrsidea* and 133 *Brueelia* individuals. *Myrsidea* occurred on all islands whereas *Brueelia* lice were only found in samples from Española, Gardner-by-Española, Gardner-by-Floreana, Marchena, Rábida and St. Fé (Fig. 2d). *Myrsidea* showed a prevalence of 95.4% and the mean number of *Myrsidea* lice in the plumage of a mockingbird was 7.6 (SD 6.5; range 0 – 35). On the six islands where *Brueelia* occurred, mean infestation of *Brueelia* was 1.26 (SD 1.7; range 0 – 10), and prevalence was 41%. Because parasite abundance has been shown to correlate positively with island size in Darwin's finches (Lindstrom et al. 2004), we also tested for an effect of island size on mean feather louse load in the different populations but found no significant correlation ($r^2 = 0.21$, $p = 0.12$).

Genotyping and genetic diversity estimates

We genotyped blood samples from a total of 400 individuals (Table 1). Successful genotypes were obtained for all individuals and loci except for one individual, which could not be genotyped at one single locus. All 26 loci were in Hardy-Weinberg equilibrium in all populations except in four cases: MpAAT96 and Nes03 in Gardner-by-Floreana, Nes04 in Santiago and Nes22 in Marchena which showed an excess of homozygotes. Genotypic disequilibrium was only detected for two locus pairs in one single population (MpAAT96 x

Nes03 and Nes05 x Nes01 in Gardner-by-Floreana). The mean number of alleles per locus was 9.5 and ranged from 3 to 18 alleles. The mean number of alleles per population across all 26 loci was 91 (range: 32-154, Table 1), mean allelic richness was 3 (range: 1.18-4.69, Table 1) and mean H_e 0.4 (range: 0.06-0.65, Table 1).

Population-specific F_{st} values were very variable and high in some populations, ranging from 0.1 for the least inbred population on St. Cruz to 0.9 for the most inbred population on Champion (Table 1). As expected, larger islands/populations showed lower F_{st} values than smaller islands ($r^2 = 0.68$, $p < 0.001$; Fig. 3). Population-specific F_{st} correlated significantly ($r^2 = 0.52$, $p = 0.006$) with temporal F_{st} . However, as the two F_{st} values represent two different measures of inbreeding, i.e. long-term versus short-term inbreeding, they were both kept in the main analyses.

Inbreeding, immunocompetence and ectoparasites

The four different response variables did not correlate (all p-values > 0.12 and r^2 -values < 0.23), except for lysis and number of feather lice which correlated positively at the population level ($r^2 = 0.36$, $p = 0.023$), but not at the individual level ($r^2 = 0.001$, $p = 0.78$). On the four islands that were sampled in all three years to account for year effects, variation between years in the four response variables did not differ significantly (all p-values > 0.07).

We found no significant effect of population-specific or temporal F_{st} on agglutination or lysis score, H:L ratio or number of feather lice (Table 2 & Fig. 2a-d). Hence, contrary to our prediction, we detected no evidence for either inbreeding estimates to affect any of our immunocompetence estimates or ectoparasite load. These results did not change when we only included the blood-sucking *Myrsidea* lice in our model, and hence omitted *Brueelia* (p-values > 0.47), or added H_e as an additional fixed effect (all p-values for $H_e > 0.22$, data not shown). However, the statistical power of the analyses was low, as reflected by the large confidence intervals of the parameter estimates (Table 2; see Hoenig and Heisey 2001 for interpretation of confidence intervals in the context of statistical power).

DISCUSSION

Inbreeding, immunocompetence and ectoparasites

Several studies support a link between inbreeding or reduced genetic diversity and susceptibility to pathogens (e.g. Coltman et al. 1999, Acevedo-Whitehouse et al. 2003, MacDougall-Shackleton et al. 2005, Cassinello et al. 2001) or immunocompetence (e.g. Whiteman et al. 2006, Reid et al. 2003, Hawley et al. 2005). Inbreeding may affect the immune system at many different levels, such as through deleterious mutations (that could lead to non-functional or less efficient proteins or cells) or a loss of heterozygous advantage, or by decreasing immune protein variation. Because decreased lymphocyte numbers (i.e. increased H:L ratio) may signal immunosuppression and because immune-compromised birds may be more affected by parasites, we predicted a positive relationship between levels of inbreeding and H:L ratio as well as feather louse load. We predicted a negative relationship with agglutination and lysis titres as more inbred birds may display less variable antibodies or deleterious mutations could affect the complement enzyme cascade. However, we detected no significant relationships between the different immune and ectoparasite estimates and either of our two inbreeding estimates (Fig. 2 & Table 2). The lack of such correlations is not uncommon, and there are several possible explanations for it.

The innate immune traits and ectoparasite load investigated here may indeed not be affected by inbreeding in the Galápagos mockingbirds. In a recent review on heterozygosity-fitness correlations in animal populations, 24% of the correlations were non-significant (Chapman et al. 2009), indicating that such results are common. Although positive heterozygosity-fitness correlations have been widely reported (for reviews see e.g. Crnokrak and Roff 1999, Keller and Waller 2002, Reed and Frankham 2003), some fitness traits, such as parasite resistance, are in many cases unrelated to genetic diversity (Poulin et al. 2000a,

Poulin et al. 2000b, Hedrick et al. 2001, Cote et al. 2005, Coltman and Slate 2003, Stevens et al. 1997, Taylor and Jamieson 2007). Rather than being associated with inbreeding or heterozygosity *per se*, genetic variation in immune traits or resistance to parasites could be associated with specific alleles (Spielman et al. 2004) such as MHC (Eizaguirre et al. 2009, Bonneaud et al. 2005). Thus, a candidate-gene approach might be more fruitful for future research to detect possible gene-resistance correlations in mockingbirds.

Alternatively, if different measures of immunity are traded-off against each other, possible correlations between inbreeding and e.g. adaptive immune response would have gone undetected in this study. If inbreeding impairs e.g. specific antibody response, there could be a shift away from adaptive immunity towards greater reliance on innate immunity, possibly reflected by the high NAb and complement enzyme levels found here (also see discussion below). H:L ratios did not seem to be affected by either of the two inbreeding measures here (Table 2) even though WBC levels have been shown to be under genetic control in other organisms (e.g. Mahaney et al. 2005, Sun et al. 2008). H:L ratios are altered by inflammation or stress and their levels therefore also reflect an individual's history of pathogen exposure. Such variation in H:L ratios may have obscured our ability to detect a genetic relationship since we could not account for the individuals' history of infection with this dataset.

Ectoparasites are only a subset of the parasites that infest a host, and as we did not examine other parasites such as endoparasites which are more likely to respond to subtle differences in the immune response (Wakelin 1996), we cannot rule out that inbreeding may have an effect on other parasites or pathogens. However, a recent health survey did not detect any blood parasites in *Mimus trifasciatus* (Deem et al. 2011).

The Galápagos Islands are exposed to strong environmental variation and pathogen pressure is known to fluctuate in such changing environments (Salam et al. 2009, Moyer et al. 2002). Therefore, mockingbirds might have adapted their innate immune systems to this wide range of conditions and purged some of their genetic load associated with immune traits

(Crnokrak and Barrett 2002) resulting in the true absence of a relationship between inbreeding and immunocompetence (e.g. Visscher et al. 2001, Wiehn et al. 2002). If purging of recessive deleterious alleles obscured a potential heterozygosity-fitness relationship (Reed and Frankham 2003, Wiehn et al. 2002), we may have expected that populations with relatively recent, fast inbreeding and strong genetic drift, such as the two small, endangered *M. trifasciatus* populations (Hoeck et al. 2010a), would show lower immunocompetence or more ectoparasites than populations with a long history of low genetic diversity (such as the populations on St. Fé or Española, Hoeck et al. 2010b). We did not detect any such pattern for the immune components measured here (Table 2 & Fig. 2), suggesting that purging is not a very likely explanation for our findings.

Alternatively, island systems, such as the Galápagos, are relatively pathogen-poor environments (McCallum and Dobson 1995, Frankham 1997, Wikelski et al. 2004) where pathogen pressure may not be sufficient for inbreeding to show a detectable effect on immunocompetence or where such effects could only become relevant under stressful environmental conditions (Arcese 2003, Nelson et al. 2002) when immune responses need to be traded-off against other traits (e.g. Hasselquist et al. 1999, Nelson et al. 2002, Zuk and Stoehr 2002). The fact that others have found evidence for a relationship between genetic diversity and pathogen load in other Galápagos birds (Whiteman et al., 2006) makes this explanation less likely. Also, lower pathogen pressure will not necessarily mask inbreeding depression in measures of innate immunity.

Although our sample size was larger than in comparable studies in Galápagos which did detect significant effects of genetic diversity or population size on immune parameters (Whiteman et al. 2006, Lindstrom et al. 2004), it is possible that some of the variation in ectoparasite infestation or immune estimates we observed is due to unaccounted environmental variability (rainfall, temperature, food availability etc.), even though we detected no significant differences in parameter estimates between years for the four islands that were sampled repeatedly. However, if such environmental effects are large in

comparison to the heterozygosity-associated signal, then the signal will be difficult to detect statistically.

The large confidence intervals associated with our estimates (Table 2) make it clear that, while we have no evidence for an association between genetic diversity and immune function, we also have no evidence that in fact such a relationship is absent. Furthermore, severe parasite infestation or disease might lead to mortality or reduced mobility and, hence, seriously sick birds might have gone undetected in our study given our capture methods. Also, we would not have detected selection that occurs early in life because we did not sample any nestlings or fledglings.

Immunocompetence measures

We did not detect any correlation between NAb levels and complement-mediated lysis or leucocyte numbers, supporting the general idea that they serve as independent measures of immunocompetence (Mendes et al. 2006, Matson et al. 2005). However, the low repeatability detected for lysis and especially agglutination cautions against such a statement. Such a low repeatability is surprising as the hemolysis-hemagglutination test has previously been shown to be a highly repeatable assay (Matson et al. 2005). We do not know what caused the low repeatability in our study. Our finding highlights the need for an independent assessment of the repeatability of this method in each individual study, something that is oftentimes omitted or not discussed.

The mean levels of agglutination in the mockingbirds (Table 1) were higher or at the high end of recently reported values from a range of birds (Matson 2006, Parejo and Silva 2009, Matson et al. 2005, Mendes et al. 2006), suggesting that Galápagos mockingbirds have high levels of natural antibodies. NAb levels are comparable to the ones found in another Galápagos endemic, the Galápagos Hawk (Whiteman et al. 2006). However, while the hawks showed no lysis reactions, all but one mockingbird population showed lysis (Table 2). The high NAb and relatively high complement enzyme titres found here could indicate that

the mockingbirds are equipped with a strong first line of defence. A shift in the immune defence strategy towards innate as opposed to acquired immune responses has previously been suggested in a study that detected increased innate defences of insular birds in comparison to their continental relatives (Matson 2006).

Conclusions

We investigated whether inbreeding at the population level affects immunocompetence in Galápagos mockingbirds, including short- and long-term inbreeding measured with recent and historic samples from populations of different sizes and inbreeding levels and examining three different measures of innate immunocompetence. We found no effects of short-term or long-term inbreeding on innate immunity or ectoparasite load. Unfortunately, despite an extensive data set our statistical power was low. The relatively high levels of agglutination and lysis detected here suggest that the mockingbirds may have a strong first line of defence against invading pathogens. However, as measures of agglutination and lysis showed very low repeatabilities, these conclusions remain tentative. Based on the results from this study, we are unable to predict how the mockingbirds would cope with new, introduced pathogens. This stresses the importance of future avian health surveys and disease control in the Galápagos.

The two endangered Floreana mockingbird (*M. trifasciatus*) populations did not show lower levels of natural antibodies, complement enzymes or higher heterophil-lymphocyte ratios than the other mockingbird populations investigated here. This suggests that *M. trifasciatus* does not suffer from immunosuppression in comparison to the more outbred and more widely distributed mockingbird species. In a recent study on the Floreana mockingbird, however, significant differences were found in different health parameters between the two satellite populations, suggesting that the genetically more diverse population on Gardner Island is clinically healthier than the smaller and more inbred Champion population (Deem et

al. 2011). Whether this results in lower fitness and survival in the Champion in comparison to the Gardner population remains subject to further investigations.

ACKNOWLEDGEMENTS

Thanks to the Galápagos National Park Service and the Charles Darwin Research Station, especially Felipe Cruz, Marilyn Cruz and Sonia Cisneros for support in Galápagos, and to Iris Biebach, Herbert Biebach, Wilson Cabrera, Stefan Henrich, María Fernanda Holguín, Michael Janssen, Karla Muñoz and Michèle Wegmann for assistance in the field. Thomas Bucher, Peter Wandeler and Andrea Gubler provided invaluable help in the lab. Kevin Matson and Richard Hoop kindly advised on the hemolysis assay and Elisabeth Rogg provided help with the identification of leucocyte types. Erik Postma, Barbara Tschirren, Iris Biebach and Simone Härri gave very useful inputs on the analyses. This research was funded by the Zoological Museum of the University of Zurich, the Forschungskredit of the University of Zurich, the Balzan Foundation (Balzan Prize 2005 awarded to Peter and Rosemary Grant), and the Basler Stiftung für biologische Forschung.

535 **REFERENCES**

- 536 ACEVEDO-WHITEHOUSE, K., GULLAND, F., GREIG, D. and AMOS, W. 2003. Disease
537 susceptibility in California sea lions. *-Nature*, **422**: 35-35.
- 538 ALLENDORF, F. W. and LUIKART, G. 2007. *Conservation and the Genetics of*
539 *Populations*, Oxford, UK, Blackwell Publishing.
- 540 APANIUS, V., YORINKS, N., BERMINGHAM, E. and RICKLEFS, R. E. 2000. Island and
541 taxon effects in parasitism and resistance of Lesser Antillean birds. *-Ecology*, **81**:
542 1959-1969.
- 543 ARCESE, P. 2003. Individual quality, environment, and conservation. In: FESTA-
544 BIANCHET, M. and APOLLONIO, M. (eds.) *Animal Behavior and Wildlife*
545 *Conservation*. Island Press, Washington, D.C., pp. 271-295.
- 546 BALLOUX, F., AMOS, W. and COULSON, T. 2004. Does heterozygosity estimate
547 inbreeding in real populations? *-Molecular Ecology*, **13**: 3021-3031.
- 548 BELPERRON, A. A. and BOCKENSTEDT, L. K. 2001. Natural antibody affects survival of
549 the spirochete *Borrelia burgdorferi* within feeding ticks. *-Infection and Immunity*, **69**:
550 6456-6462.
- 551 BIEBACH, I. and KELLER, L. F. 2010. Inbreeding in reintroduced populations: the effects of
552 early reintroduction history and contemporary processes. *-Conservation Genetics*, **11**:
553 527-538.
- 554 BONNEAUD, C., RICHARD, M., FAIVRE, B., WESTERDAHL, H. and SORCI, G. 2005.
555 An Mhc class I allele associated to the expression of T-dependent immune response in
556 the house sparrow. *-Immunogenetics*, **57**: 782-789.
- 557 BOOTH, D. T., CLAYTON, D. H. and BLOCK, B. A. 1993. Experimental Demonstration of
558 the Energetic Cost of Parasitism in Free-Ranging Hosts. *-Proceedings of the Royal*
559 *Society of London Series B-Biological Sciences*, **253**: 125-129.
- 560 BOX, G. E. P. and COX, D. R. 1964. An Analysis Of Transformations. *-Journal Of The*
561 *Royal Statistical Society Series B-Statistical Methodology*, **26**: 211-252.
- 562 CARROLL, M. C. and PRODEUS, A. P. 1998. Linkages of innate and adaptive immunity. -
563 *Current Opinion in Immunology*, **10**: 36-40.
- 564 CASSINELLO, J., GOMENDIO, M. and ROLDAN, E. R. S. 2001. Relationship between
565 coefficient of inbreeding and parasite burden in endangered gazelles. *-Conservation*
566 *Biology*, **15**: 1171-1174.
- 567 CHAPMAN, J. R., NAKAGAWA, S., COLTMAN, D. W., SLATE, J. and SHELDON, B. C.
568 2009. A quantitative review of heterozygosity-fitness correlations in animal
569 populations. *-Molecular Ecology*, **18**: 2746-2765.
- 570 CHENG, S. and LAMONT, S. J. 1988. Genetic-Analysis of Immunocompetence Measures in
571 a White Leghorn Chicken Line. *-Poultry Science*, **67**: 989-995.
- 572 CIOFI, C., BEAUMONT, M. A., SWINGLAND, I. R. and BRUFORD, M. W. 1999. Genetic
573 divergence and units for conservation in the Komodo dragon *Varanus komodoensis*. -
574 *Proceedings of the Royal Society of London Series B-Biological Sciences*, **266**: 2269-
575 2274.
- 576 CLAYTON, D. H., LEE, P. L. M., TOMPKINS, D. M. and BRODIE, E. D. 1999. Reciprocal
577 natural selection on host-parasite phenotypes. *-American Naturalist*, **154**: 261-270.
- 578 COLTMAN, D. W., PILKINGTON, J. G., SMITH, J. A. and PEMBERTON, J. M. 1999.
579 Parasite-mediated selection against inbred Soay sheep in a free-living, island
580 population. *-Evolution*, **53**: 1259-1267.
- 581 COLTMAN, D. W. and SLATE, J. 2003. Microsatellite measures of inbreeding: A meta-
582 analysis. *-Evolution*, **57**: 971-983.

- COTE, S. D., STIEN, A., IRVINE, R. J., DALLAS, J. F., MARSHALL, F., HALVORSEN, O., LANGVATN, R. and ALBON, S. D. 2005. Resistance to abomasal nematodes and individual genetic variability in reindeer. *-Molecular Ecology*, **14**: 4159-4168.
- CRNOKRAK, P. and BARRETT, S. C. H. 2002. Perspective: Purging the genetic load: A review of the experimental evidence. *-Evolution*, **56**: 2347-2358.
- CRNOKRAK, P. and ROFF, D. A. 1999. Inbreeding depression in the wild. *-Heredity*, **83**: 260-270.
- DAVIS, A. K., MANEY, D. L. and MAERZ, J. C. 2008. The use of leukocyte profiles to measure stress in vertebrates: a review for ecologists. *-Functional Ecology*, **22**: 760-772.
- DEEM, S. L., PARKER, P. G., CRUZ, M. B., MERKEL, J. and HOECK, P. E. A. 2011. Comparison of blood values and health status of Floreana Mockingbirds (*Mimus trifasciatus*) on the islands of Champion and Gardner-by-Floreana, Galapagos Islands - *Journal of Wildlife Diseases*, **47**: 94-106.
- DEWOODY, Y. D. and DEWOODY, J. A. 2005. On the estimation of genome-wide heterozygosity using molecular markers. *-Journal of Heredity*, **96**: 85-88.
- EIZAGUIRRE, C., YEATES, S. E., LENZ, T. L., KALBE, M. and MILINSKI, M. 2009. MHC-based mate choice combines good genes and maintenance of MHC polymorphism. *-Molecular Ecology*, **18**: 3316-3329.
- FRANKHAM, R. 1997. Do island populations have less genetic variation than mainland populations? *-Heredity*, **78**: 311-327.
- GERLOFF, C. U., OTTMER, B. K. and SCHMID-HEMPEL, P. 2003. Effects of inbreeding on immune response and body size in a social insect, *Bombus terrestris*. *-Functional Ecology*, **17**: 582-589.
- GIESE, A. R. and HEDRICK, P. W. 2003. Genetic variation and resistance to a bacterial infection in the endangered Gila topminnow. *-Animal Conservation*, **6**: 369-377.
- GOTTDENKER, N. L., WALSH, T., VARGAS, H., MERKEL, J., JIMENEZ, G. U., MILLER, R. E., DAILEY, M. and PARKER, P. G. 2005. Assessing the risks of introduced chickens and their pathogens to native birds in the Galapagos Archipelago. *-Biological Conservation*, **126**: 429-439.
- GOUDET, J. 2001. FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3). Lausanne University, Lausanne, Switzerland.
- GRIFFITHS, R., DOUBLE, M. C., ORR, K. and DAWSON, R. J. G. 1998. A DNA test to sex most birds. *-Molecular Ecology*, **7**: 1071-1075.
- GROSS, W. B. and SIEGEL, H. S. 1983. Evaluation of the Heterophil Lymphocyte Ratio as a Measure of Stress in Chickens. *-Avian Diseases*, **27**: 972-979.
- HARRIS, M. P. 1974. *The birds of Galapagos*, New York, Houghton-Mifflin.
- HASSELQUIST, D., MARSH, J. A., SHERMAN, P. W. and WINGFIELD, J. C. 1999. Is avian humoral immunocompetence suppressed by testosterone? *-Behavioral Ecology and Sociobiology*, **45**: 167-175.
- HAWLEY, D. M., DHONDT, K. V., DOBSON, A. P., GRODIO, J. L., HOCHACHKA, W. M., LEY, D. H., OSNAS, E. E., SCHAT, K. A. and DHONDT, A. A. 2010. Common garden experiment reveals pathogen isolate but no host genetic diversity effect on the dynamics of an emerging wildlife disease. *-Journal of Evolutionary Biology*, **23**: 1680-1688.
- HAWLEY, D. M., SYDENSTRICKER, K. V., KOLLIAS, G. V. and DHONDT, A. A. 2005. Genetic diversity predicts pathogen resistance and cell-mediated immunocompetence in house finches. *-Biology Letters*, **1**: 326-329.
- HEDRICK, P. W., KIM, T. J. and PARKER, K. M. 2001. Parasite resistance and genetic variation in the endangered Gila topminnow. *-Animal Conservation*, **4**: 103-109.

- HOECK, P. E. A., BEAUMONT, M. A., JAMES, K. E., GRANT, B. R., GRANT, P. R. and KELLER, L. F. 2010a. Saving Darwin's muse: evolutionary genetics for the recovery of the Floreana mockingbird. *-Biology Letters*, **6**: 212-215.
- HOECK, P. E. A., BOLLMER, J. L., PARKER, P. G. and KELLER, L. F. 2010b. Differentiation with drift: a spatio-temporal genetic analysis of Galapagos mockingbird populations (*Mimus* spp.). *-Philosophical Transactions of the Royal Society B-Biological Sciences*, **365**: 1127-1138.
- HOECK, P. E. A., BUCHER, T. B., WANDELER, P. and KELLER, L. F. 2009. Microsatellite primers for the four Galapagos mockingbird species (*Mimus parvulus*, *Mimus macdonaldi*, *Mimus melanotis* and *Mimus trifasciatus*). *-Molecular Ecology Resources*, **9**: 1538-1541.
- HOENIG, J. M. and HEISEY, D. M. 2001. The abuse of power: The pervasive fallacy of power calculations for data analysis. *-American Statistician*, **55**: 19-24.
- KELLER, L. F. and WALLER, D. M. 2002. Inbreeding effects in wild populations. *-Trends in Ecology & Evolution*, **17**: 230-241.
- KOHLER, H., BAYRY, J., NICOLETTI, A. and KAVERI, S. V. 2003. Natural autoantibodies as tools to predict the outcome of immune response? *-Scandinavian Journal of Immunology*, **58**: 285-289.
- LAMMERS, A., KLOMP, M. E. V., NIEUWLAND, M. G. B., SAVELKOUL, H. F. J. and PARMENTIER, H. K. 2004. Adoptive transfer of natural antibodies to non-immunized chickens affects subsequent antigen-specific humoral and cellular immune responses. *-Developmental and Comparative Immunology*, **28**: 51-60.
- LESSELLS, C. M. and BOAG, P. T. 1987. Unrepeatable repeatabilities - a common mistake. *-Auk*, **104**: 116-121.
- LINDSTROM, K. M., FOUFOPOULOS, J., PARN, H. and WIKELSKI, M. 2004. Immunological investments reflect parasite abundance in island populations of Darwin's finches. *-Proceedings of the Royal Society of London Series B-Biological Sciences*, **271**: 1513-1519.
- LJUNGQVIST, M., AKESSON, M. and HANSSON, B. 2010. Do microsatellites reflect genome-wide genetic diversity in natural populations? A comment on Vali et al. (2008). *-Molecular Ecology*, **19**: 851-855.
- LUONG, L. T., HEATH, B. D. and POLAK, M. 2007. Host inbreeding increases susceptibility to ectoparasitism. *-Journal of Evolutionary Biology*, **20**: 79-86.
- MACDOUGALL-SHACKLETON, E. A., DERRYBERRY, E. P., FOUFOPOULOS, J., DOBSON, A. P. and HAHN, T. P. 2005. Parasite-mediated heterozygote advantage in an outbred songbird population. *-Biology Letters*, **1**: 105-107.
- MAHANEY, M. C., BRUGNARA, C., LEASE, L. R. and PLATT, O. S. 2005. Genetic influences on peripheral blood cell counts: a study in baboons. *-Blood*, **106**: 1210-1214.
- MARSHALL, A. G. 1981. *The Ecology of Ectoparasitic Insects*, London, Academic Press.
- MATSON, K. D. 2006. Are there differences in immune function between continental and insular birds? *-Proceedings of the Royal Society B-Biological Sciences*, **273**: 2267-2274.
- MATSON, K. D., RICKLEFS, R. E. and KLASING, K. C. 2005. A hemolysis-hemagglutination assay for characterizing constitutive innate humoral immunity in wild and domestic birds. *-Developmental and Comparative Immunology*, **29**: 275-286.
- MAXWELL, M. H. 1993. Avian Blood Leukocyte Responses to Stress. *-Worlds Poultry Science Journal*, **49**: 34-43.
- MCCALLUM, H. and DOBSON, A. 1995. Detecting Disease and Parasite Threats to Endangered Species and Ecosystems. *-Trends in Ecology & Evolution*, **10**: 190-194.

- MENDES, L., PIERSMA, T., HASSELQUIST, D., MATSON, K. D. and RICKLEFS, R. E. 2006. Variation in the innate and acquired arms of the immune system among five shorebird species. -*Journal of Experimental Biology*, **209**: 284-291.
- MILLER, H. C. and LAMBERT, D. M. 2004. Genetic drift outweighs balancing selection in shaping post-bottleneck major histocompatibility complex variation in New Zealand robins (Petroicidae). -*Molecular Ecology*, **13**: 3709-3721.
- MOLLER, A. P. and ROZSA, L. 2005. Parasite biodiversity and host defenses: chewing lice and immune response of their avian hosts. -*Oecologia*, **142**: 169-176.
- MOYER, B. R., DROWN, D. M. and CLAYTON, D. H. 2002. Low humidity reduces ectoparasite pressure: implications for host life history evolution. -*Oikos*, **97**: 223-228.
- NELSON, R. J., DEMAS, G. E., KLEIN, S. L. and KRIEGSFELD, L. J. 2002. *Seasonal Patterns of Stress, Immune Function and Disease*, Cambridge, Cambridge University Press.
- NORRIS, K. and EVANS, M. R. 2000. Ecological immunology: life history trade-offs and immune defense in birds. -*Behavioral Ecology*, **11**: 19-26.
- OCHSENBEIN, A. F. and ZINKERNAGEL, R. M. 2000. Natural antibodies and complement link innate and acquired immunity. -*Immunology Today*, **21**: 624-630.
- PAREJO, D. and SILVA, N. 2009. Immunity and fitness in a wild population of Eurasian kestrels *Falco tinnunculus*. -*Naturwissenschaften*, **96**: 1193-1202.
- PARKER, P. G., WHITEMAN, N. K. and MILLER, R. E. 2006. Conservation medicine on the galapagos islands: Partnerships among behavioral, population, and veterinary scientists. -*Auk*, **123**: 625-638.
- PARMENTIER, H. K., LAMMERS, A., HOEKMAN, J. J., REILINGH, G. D., ZAAANEN, I. T. A. and SAVELKOUL, H. F. J. 2004. Different levels of natural antibodies in chickens divergently selected for specific antibody responses. -*Developmental and Comparative Immunology*, **28**: 39-49.
- PEARMAN, P. B. and GARNER, T. W. J. 2005. Susceptibility of Italian agile frog populations to an emerging strain of Ranavirus parallels population genetic diversity. -*Ecology Letters*, **8**: 401-408.
- POULIN, R., MARSHALL, L. J. and SPENCER, H. G. 2000a. Genetic variation and prevalence of blood parasites do not correlate among bird species. -*Journal of Zoology*, **252**: 381-388.
- POULIN, R., MARSHALL, L. J. and SPENCER, H. G. 2000b. Metazoan parasite species richness and genetic variation among freshwater fish species: cause or consequence? -*International Journal for Parasitology*, **30**: 697-703.
- PUURTINEN, M., HYTONEN, M., KNOTT, K. E., TASKINEN, J., NISSINEN, K. and KAITALA, V. 2004. The effects of mating system and genetic variability on susceptibility to trematode parasites in a freshwater snail, *Lymnaea stagnalis*. -*Evolution*, **58**: 2747-2753.
- REED, D. H. 2004. Extinction risk in fragmented habitats. -*Animal Conservation*, **7**: 181-191.
- REED, D. H. and FRANKHAM, R. 2003. Correlation between fitness and genetic diversity. -*Conservation Biology*, **17**: 230-237.
- REID, J. M., ARCESE, P. and KELLER, L. F. 2003. Inbreeding depresses immune response in song sparrows (*Melospiza melodia*): direct and inter-generational effects. -*Proceedings of the Royal Society of London Series B-Biological Sciences*, **270**: 2151-2157.
- REID, J. M., ARCESE, P., KELLER, L. F., ELLIOTT, K. H., SAMPSON, L. and HASSELQUIST, D. 2007. Inbreeding effects on immune response in free-living song sparrows (*Melospiza melodia*). -*Proceedings of the Royal Society B-Biological Sciences*, **274**: 697-706.

- RICKLEFS, R. E. and SHELDON, K. S. 2007. Malaria prevalence and white-blood-cell response to infection in a tropical and in a temperate thrush. *-Auk*, **124**: 1254-1266.
- ROSS-GILLESPIE, A., O'RIAIN, M. J. and KELLER, L. F. 2007. Viral epizootic reveals inbreeding depression in a habitually inbreeding mammal. *-Evolution*, **61**: 2268-2273.
- SALAM, S. T., MIR, M. S. and KHAN, A. R. 2009. Prevalence and seasonal variation of ectoparasite load in free-range chicken of Kashmir valley. *-Tropical Animal Health and Production*, **41**: 1371-1376.
- SANDLAND, G. J., WETHINGTON, A. R., FOSTER, A. V. and MINCHELLA, D. J. 2009. Effects of host outcrossing on the interaction between an aquatic snail and its locally adapted parasite. *-Parasitology Research*, **105**: 555-561.
- SARKER, N., TSUDZUKI, M., NISHIBORI, M., YASUE, H. and YAMAMOTO, Y. 2000. Cell-mediated and humoral immunity and phagocytic ability in chicken lines divergently selected for serum immunoglobulin M and G levels. *-Poultry Science*, **79**: 1705-1709.
- SMITH, K. F., ACEVEDO-WHITEHOUSE, K. and PEDERSEN, A. B. 2009. The role of infectious diseases in biological conservation. *-Animal Conservation*, **12**: 1-12.
- SOOS, C., PADILLA, L., IGLESIAS, A., GOTTDENKER, N., BEDON, M. C., RIOS, A. and PARKER, P. G. 2008. Comparison of pathogens in broiler and backyard chickens on the Galapagos Islands: Implications for transmission to wildlife. *-Auk*, **125**: 445-455.
- SPIELMAN, D., BROOK, B. W., BRISCOE, D. A. and FRANKHAM, R. 2004. Does inbreeding and loss of genetic diversity decrease disease resistance? *-Conservation Genetics*, **5**: 439-448.
- STEFKA, J., HOECK, P. E. A., KELLER, L. F. and SMITH, V. S. 2011. A hitchhikers guide to the Galapagos: co-phylogeography of Galapagos mockingbirds and their parasites. *-Bmc Evolutionary Biology*, **11**.
- STEVENS, L., YAN, G. Y. and PRAY, L. A. 1997. Consequences of inbreeding on invertebrate host susceptibility to parasitic infection. *-Evolution*, **51**: 2032-2039.
- SUN, S. F., PAN, Q. Z., HUI, X., ZHANG, B. L., WU, H. M., LI, H., XU, W., ZHANG, Q., LI, J. Y., DENG, X. M., CHEN, J. W., LIAN, Z. X. and LI, N. 2008. Stronger In Vitro Phagocytosis by Monocytes-Macrophages Is Indicative of Greater Pathogen Clearance and Antibody Levels In Vivo. *-Poultry Science*, **87**: 1725-1733.
- TAYLOR, S. S. and JAMIESON, I. G. 2007. Factors affecting the survival of founding individuals in translocated New Zealand Saddlebacks *Philesturnus carunculatus*. *-Ibis*, **149**: 783-791.
- TOMPKINS, D. M., MITCHELL, R. A. and BRYANT, D. M. 2006. Hybridization increases measures of innate and cell-mediated immunity in an endangered bird species. *-Journal of Animal Ecology*, **75**: 559-564.
- VAN RIPER, C., VAN RIPER, S. G., GOFF, M. L. and LAIRD, M. 1986. The Epizootiology and Ecological Significance of Malaria in Hawaiian Land Birds. *-Ecological Monographs*, **56**: 327-344.
- VAN RIPER, C., VAN RIPER, S. G. and HANSEN, W. R. 2002. Epizootiology and effect of avian pox on Hawaiian forest birds. *-Auk*, **119**: 929-942.
- VISSCHER, P. M., SMITH, D., HALL, S. J. G. and WILLIAMS, J. A. 2001. A viable herd of genetically uniform cattle - Deleterious alleles seem to have been purged in a feral strain of inbred cows. *-Nature*, **409**: 303-303.
- WAKELIN, D. 1996. *Immunity to Parasites. How Parasitic Infections are Controlled*, Cambridge, Cambridge University Press.

- WALTHER, B. A. and CLAYTON, D. H. 1997. Dust-ruffling: A simple method for quantifying ectoparasite loads of live birds. *-Journal of Field Ornithology*, **68**: 509-518.
- WHITEMAN, N. K., MATSON, K. D., BOLLMER, J. L. and PARKER, P. G. 2006. Disease ecology in the Galapagos Hawk (*Buteo galapagoensis*): host genetic diversity, parasite load and natural antibodies. *-Proceedings of the Royal Society B-Biological Sciences*, **273**: 797-804.
- WIEHN, J., KOPP, K., REZZONICO, S., KARTTUNEN, S. and JOKELA, J. 2002. Family-level covariation between parasite resistance and mating system in a hermaphroditic freshwater snail. *-Evolution*, **56**: 1454-1461.
- WIKEL, S. K. and ALARCON-CHAIDEZ, F. J. 2001. Progress toward molecular characterization of ectoparasite modulation of host immunity. *-Veterinary Parasitology*, **101**: 275-287.
- WIKELSKI, M., FOUFOPOULOS, J., VARGAS, H. and SNELL, H. 2004. Galapagos Birds and Diseases: Invasive Pathogens as Threats for Island Species. *-Ecology and Society*, **9**.
- WRIGHT, S. 1969. *Evolution and the Genetics of Populations. The Theory of Gene Frequencies.*, Chicago, University of Chicago Press.
- ZOUALI, M. 2001. Natural Antibodies. *Encyclopedia of Life Sciences*. John Wiley & Sons, Ltd.
- ZUK, M. and JOHNSEN, T. S. 1998. Seasonal changes in the relationship between ornamentation and immune response in red jungle fowl. *-Proceedings of the Royal Society of London Series B-Biological Sciences*, **265**: 1631-1635.
- ZUK, M. and STOEHR, A. M. 2002. Immune defense and host life history. *-American Naturalist*, **160**: S9-S22.

FIGURE LEGENDS

Figure 1:

Distribution of the four different mockingbird species (*Mimus* spp.) in the Galápagos archipelago with circles indicating the sites of sample collection. On St. Cruz (*PtA* and *GP*) and San Cristóbal (*Sur* and *Nor*) samples were obtained from two different locations.

Figure 2:

The three estimates of immunocompetence and feather louse numbers shown as a function of population-specific F_{st} , a measure of inbreeding, with values of F_{st} increasing from left to right (real numbers are given in Table 1 and Figure 3). To improve visual representation, points that overlapped (on the x-axis) were slightly moved. Bars represent the standard errors of the means. a) The mean heterophil-lymphocyte (H:L) ratio of each population. b) Mean lysis and c) mean agglutination scores for each population, with values corrected for the chicken control sample. d) The mean number of *Myrsidea* (closed circles) and *Brueelia* (open circles) lice per population. *Brueelia* did not occur in all populations. The total mean number of feather lice per population is shown in Table 1.

Figure 3:

Population-specific F_{st} as a function of the natural logarithm of island size (in ha; actual values are given in Table 1) with island size decreasing from left to right. Points that overlapped on the x-axis were slightly moved to improve visual representation. The two F_{st} estimates for St. Cruz and San Cristóbal represent the two different sampling sites on these islands. Bars show the minimum and maximum estimates for F_{st} as calculated in 2mod (Ciofi et al. 1999).

836 TABLES

837

838 Table 1:

839 Populations studied with island size (log-scale, in hectares), estimates of genetic diversity
 840 and immunocompetence and the respective sample sizes (N, number of mockingbird
 841 samples). Measures of genetic diversity are based on 26 microsatellite loci: NA: number of
 842 alleles, AR: allelic richness, He: expected heterozygosity, pop. Fst: population-specific Fst as
 843 calculated in 2mod (Ciofi et al. 1999), temp. Fst: temporal Fst as calculated in Hoeck et al.
 844 (2010b). Mean lysis and agglutination scores are given (for chicken control corrected values
 845 see Fig. 2b & c), as well as mean heterophil-lymphocyte (H:L) ratio and mean number of total
 846 feather lice encountered per individual in each population.

847

Island	island size (ln; ha)	genetic diversity				inbreeding		hemolysis- hemagglutination			H:L ratio		# of feather lice	
		NA	AR	He	N	pop. Fst	temp. Fst	mean lysis score	mean agglutination score	N	mean	N	mean	N
Champion	2.25	32	1.18	0.06	48	0.90	0.28	3.27	9.33	24	0.20	62	6.83	12
Espanola	8.71	68	2.08	0.25	58	0.65	0.02	3.58	8.48	63	0.33	88	9.27	33
Gardner-by-Espanola	4.06	45	1.73	0.17	10	0.72	0.11	4.33	9.33	3	0.44	10	13.89	9
Gardner-by-Floreana	4.4	62	2.12	0.32	69	0.69	0.08	3.68	9.18	44	0.21	72	8.45	22
Isabela	13.04	128	3.77	0.48	32	0.27	0.02	2.95	9.19	16	0.21	29	8.07	14
Marchena	9.47	95	3.04	0.47	38	0.39	0.02	3.55	8.45	20	0.11	39	8.11	18
Rabida	6.21	96	3.29	0.51	21	0.32	0.05	.	9.13	15	0.32	21	12.50	10
SanCristobal-Sur	10.93	96	3.23	0.40	20	0.39	0.02	2.50	9.46	14	0.11	12	2.50	6
SanCristobal-Nor	10.93	79	2.78	0.37	17	0.47	0.02	3.42	9.31	13	0.14	17	6.67	15
Santiago	10.98	154	4.69	0.61	27	0.12	0.01	3.70	7.37	15	0.12	27	2.25	12
StCruz-Gar	11.5	133	4.37	0.63	22	0.14	0.01	3.17	8.35	17	0.23	38	8.75	16
StCruz-PtA	11.5	137	4.68	0.65	17	0.11	0.01	3.31	8.77	13	0.21	17	4.89	9
StFe	7.79	59	2.13	0.33	21	0.58	0.01	3.71	9.71	12	0.10	21	10.30	20
Total # of samples					400					269		453		196

848

849

850


851

Table 2:

GLMs for a) agglutination (n=13) and b) lysis scores (n=12), corrected for chicken control sample, c) heterophil-lymphocyte (H:L) ratio (ln-transformed, n=13), and d) number of feather lice (transformed to the power of 0.25, n=13).


	Estimate	±SE	95%CI	t-value	d.f.	p-value
a) Agglutination						
Intercept	-1.57	0.56	-2.8 – -0.3	-2.82	1,10	
population-specific Fst	2.23	1.42	-0.9 – 5.4	1.56	1,10	0.15
temporal Fst	-2.24	4.76	-13.0 – 8.5	-0.47	1,10	0.65
b) Lysis						
Intercept	-0.91	0.35	-1.7 – -0.1	-2.64	1, 9	
population-specific Fst	1.11	0.89	-0.9 – 3.1	1.25	1, 9	0.24
temporal Fst	-1.21	2.97	-7.9 – 5.5	-0.41	1, 9	0.69
c) H:L ratio						
Intercept	-2.30	0.31	-2.9 – -1.6	-7.51	1,10	
population-specific Fst	0.47	0.79	-1.3 – 2.2	0.59	1,10	0.57
temporal Fst	0.88	2.63	-5.1 – 6.8	0.33	1,10	0.75
d) Feather lice						
Intercept	1.24	0.16	0.8 – 1.6	7.77	1,10	
population-specific Fst	0.59	0.41	-0.3 – 1.5	1.45	1,10	0.18
temporal Fst	-0.75	1.37	-3.9 – 2.4	-0.55	1,10	0.60

859 **Figure 1:**
860




861
862
863
864

Figure 2:



870 **Figure 3:**
871



872